arXiv:0905.1294v1 [math.CA] 8 May 2009

A new class of numerical sequences and its applications to uniform convergence of sine series

Bogdan Szal
University of Zielona Góra,
Faculty of Mathematics, Computer Science and Econometrics,
65-516 Zielona Góra, ul. Szafrana 4a, Poland
e-mail: B.Szal @wmie.uz.zgora.pl

Abstract

In the present paper we introduce a new class of sequences called $GM(\beta, r)$, which is the generalization of a class considered by Tikhonov in [13]. Moreover, we obtained in this note sufficient and necessary conditions for uniform convergence of sine series with (β, r) −general monotone coefficients.

Keywords and phrases: Sine series, Fourier series, embedding relations, number sequences.

2000 Mathematics Subject Classification: 40A30, 42A10

1 Introduction

It is well know that there are a great number of interesting results in Fourier analysis established by assuming monotonicity of coefficients. The following classical convergence result can be found in many monographs (see [1] and [18], for example).

Theorem 1. *Suppose that* $b_n \geq b_{n+1}$ *and* $b_n \to 0$. *Then a necessary and sufficient condition for the uniform convergence of the series*

$$\sum_{n=1}^{\infty} b_n \sin nx \tag{1.1}$$

is $nb_n \to 0$.

This result have been generalized by weakening the monotone conditions of the coefficient sequences. Generally speaking, it has become an important topic how to generalize monotonicity.

Recently, Leindler [3] defined a new class of sequences named as sequences of rest bounded variation, briefly denoted by $RBVS$, i.e.,

$$RBVS=\left\{a:=(a_n)\in\mathbb{C}:\sum_{n=m}^{\infty}|a_n-a_{n+1}|\leq K(a)\,|a_m|\ \text{ for all } m\in\mathbb{N}\right\},$$

where here and throughout the paper $K(a)$ always indicates a constant only depending on a.

Denote by MS the class of monotone decreasing sequences and $CQMS$ the class of classic quasimonotone decreasing sequences ($a\in CQMS$ means that $(a_n)\in\mathbb{R}_+$ and there exists an $\alpha>0$ such that a_n/n^{α} is decreasing), then it is obvious that

$$MS\subset RBVS\cap CQMS. \tag{1.2}$$

Leindler [4] proved that the class $CQMS$ and $RBVS$ are not comparable. In [9] leindler considered the class of mean rest bounded variation sequences $MRBVS$, where

$$\begin{gathered}MRBVS=\{a:=(a_n)\in\mathbb{C}:\\ \left.\sum_{n=m}^{\infty}|a_n-a_{n+1}|\leq K(a)\,\frac{1}{m}\sum_{n\geq m/2}^{m}|a_n|\ \text{ for all } m\in\mathbb{N}\right\}.\end{gathered}$$

Further, the class of general monotone coefficients, GM, is defined as follows (see [11]):

$$GM=\left\{a:=(a_n)\in\mathbb{C}:\sum_{n=m}^{2m-1}|a_n-a_{n+1}|\leq K(a)\,|a_m|\ \text{ for all } m\in\mathbb{N}\right\}.$$

It is clear that

$$RBVS\subset MRBVS\quad\text{and}\quad RBVS\cup CQMS\subset GM. \tag{1.3}$$

Very recently, Le and Zhou [2] suggested the following new class of sequences to include GM:

$$\begin{gathered}GBVS=\{a:=(a_n)\in\mathbb{C}:\\ \left.\sum_{n=m}^{2m-1}|a_n-a_{n+1}|\leq K(a)\max_{m\leq n\leq N+m}|a_n|\ \text{ for some integer } N \text{ and all } m\in\mathbb{N}\right\}\end{gathered}$$

In [7, 11, 12, 13] was defined the class of $\beta-$general monotone sequences as follows:

Definition 1. Let $\beta := (\beta_n)$ be a nonnegative sequence. The sequence of complex numbers $a := (a_n)$ is said to be $\beta-$general monotone, or $a \in GM(\beta)$, if the relation

$$\sum_{n=m}^{2m-1} |a_n - a_{n+1}| \leq K(a)\,\beta_m$$

holds for all m.

In the paper [13] Tikhonov considered the following examples of the sequences β_n :

(1) ${}_1\beta_n = |a_n|$,

(2) ${}_2\beta_n = \sum\limits_{k=n}^{n+N} |a_k|$ for some integer N,

(3) ${}_3\beta_n = \sum\limits_{\nu=0}^{N} |a_{c^\nu n}|$ for some integers N and $c > 1$,

(4) ${}_4\beta_n = |a_n| + \sum\limits_{k=n+1}^{[cn]} \frac{|a_k|}{k}$ for some $c > 1$,

(5) ${}_5\beta_n = |a_n| + \sum\limits_{k=[n/c]}^{[cn]} \frac{|a_k|}{k}$ for some $c > 1$.

It is clear that $GM({}_1\beta) = GM$ and $GM({}_2\beta) = GBVS$. Moreover (see [13, Remark 2.1])

$$GM({}_1\beta +_2 \beta +_3 \beta +_4 \beta +_5 \beta) \equiv GM({}_5\beta). \tag{1.4}$$

The following results generalized the Chaudy - Joliffe criteria (Theorem 1) as well as its extensions (see also (1.2), (1.3) and (1.4)): [10] for $CQMS$, [3] for $RBVS$, [14] for $MRBVS$, [11] for GM, [2] for $GBVS$, [15] for particular case of $GM({}_3\beta)$ sequences, [16, 17] for $GM({}_5\beta)$, [7] and [12] for $GM(\beta)$ but only the sufficient condition and [13] for $GM(\beta)$.

We write $I_1 \ll I_2$ if there exists a positive constant K such that $I_1 \leq K I_2$.

In order to formulate our new results we define another class of sequences.

Definition 2. Let $\beta := (\beta_n)$ be a nonnegative sequence and r a natural number. The sequence of complex number $a := (a_n)$ is said to be $(\beta, r)-$general monotone, or $a \in GM(\beta, r)$, if the relation

$$\sum_{n=m}^{2m-1} |a_n - a_{n+r}| \leq K(a)\,\beta_m$$

holds for all m.

It is clear that $GM\left(\beta,1\right)\equiv GM\left(\beta\right)$. Moreover, the embedding relation between $GM\left(\beta,r\right)$ $(r>1)$ and $GM\left(\beta,1\right)$ implies from the following remark:

Remark 1. Let r be a natural number. If a nonnegative sequence $\beta:=\left(\beta_n\right)$ is such that

$$\sum_{i=0}^{r-1}\beta_{n+i}\ll\beta_n$$

for all n, then

$$GM\left(\beta,1\right)\subseteq GM\left(\beta,r\right).$$

Connecting with the relation (1.4) we shall consider the class $GM\left(\beta^{*},r\right)$ only, where

$$\beta^{*}:=\beta^{*}\left(r\right)=\sum_{k=n}^{n+r-1}\left|a_k\right|+\sum_{k=[n/c]}^{[cn]}r\frac{\left|a_k\right|}{k}\text{ for some }c>1.$$

It is clear that ${}_5\beta\equiv\beta^{*}\left(1\right)$ and $GM\left({}_5\beta\right)\equiv GM\left(\beta^{*},1\right)\subseteq GM\left(\beta^{*},r\right)$ for $r\geq 1$ (see Corollary 1).

In this note we shall present the properties of the class $GM\left(\beta^{*},r\right)$. Moreover, we generalize and extend to the class $GM\left(\beta^{*},r\right)$ the results of Tikhonov, which are included in [13].

2 Main results

We have the following results:

Theorem 2. *Let* $r_1,r_2\in\mathbb{N}$ *and* $r_1<r_2$. *If* $r_1\mid r_2$, *then* $GM\left(\beta^{*},r_1\right)\subsetneq GM\left(\beta^{*},r_2\right)$.

Corollary 1. *If* $r\in\mathbb{N}$ *and* $r>1$, *then* $GM\left({}_5\beta\right)=GM\left(\beta^{*},1\right)\subsetneq GM\left(\beta^{*},r\right)$.

Theorem 3. *Let* $r_1,r_2\in\mathbb{N}$. *If* $r_1\nmid r_2$ *and* $r_2\nmid r_1$, *then the classes* $GM\left(\beta^{*},r_1\right)$ *and* $GM\left(\beta^{*},r_2\right)$ *are not comparable.*

Theorem 4. *Let a nonnegative sequence* $\left(b_n\right)\in GM\left(\beta^{*},r\right)$, *where* $r\geq 1$. *If the series (1.1) converges uniformly (or if the series (1.1) is the Fourier series of a continuous function), then* $nb_n\rightarrow 0$ *as* $n\rightarrow\infty$.

Theorem 5. *Let a sequence* $\left(b_n\right)\in GM\left(\beta^{*},2\right)$. *If* $n\left|b_n\right|\rightarrow 0$ *as* $n\rightarrow\infty$, *then the series (1.1) converges uniformly.*

Corollary 2. *Let a nonnegative sequence* $(b_n) \in GM(\beta^*, 2)$. *Then the necessary and sufficient condition for series (1.1) to be uniformly convergent is* $nb_n \to 0$ *as* $n \to \infty$.

Remark 2. If we confine our attention to the class $GM({}_5\beta)$ then by Corollary 1 the Tikhonov result (see [13, Remark 2.2.2]) follows from Corollary 2.

Remark 3. There exist a real number x_0 and a sequence $d := (d_n) \in GM(\beta^*, 3)$, with the property $nd_n \to 0$ as $n \to \infty$, for which the series (1.1) is divergent in x_0.

Theorem 6. *Let a sequence* $(b_n) \in GM(\beta^*, r)$, *where* $r \geq 3$. *If* $n|b_n| \to 0$ *as* $n \to \infty$ *and*

$$\sum_{n=1}^{\infty} \sum_{k=1}^{[r/2]} |b_{r\cdot n+k} - b_{r\cdot n+r-k}| < \infty,$$

then the series (1.1) converges uniformly.

Remark 4. The above result is essential extension of the Tikhonov result (see [13, Theorem 2.1]). Indeed, for any $r \geq 3$ there exists a sequence $a := (a_n) \in GM(\beta^*, r)$, with the properties: $na_n \to 0$ as $n \to \infty$ and

$$\sum_{n=1}^{\infty} \sum_{k=1}^{[r/2]} |a_{r\cdot n+k} - a_{r\cdot n+r-k}| < \infty, \tag{2.1}$$

which does not belong to the class $GM(\beta^*, 2)$.

3 Lemmas

Lemma 1. *Let* $r \in \mathbb{N}$, $l \in \mathbb{Z}$ *and* $a := (a_n) \in \mathbb{C}$. *If* $x \neq \frac{2l\pi}{r}$, *then for all* n

$$\sum_{k=n}^{2n-1} a_k \sin kx = \frac{-1}{2\sin(rx/2)} \left\{ \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \cos\left(k + \frac{r}{2}\right)x \right.$$
$$\left. + \sum_{k=2n}^{2n+r-1} a_k \cos\left(k - \frac{r}{2}\right)x - \sum_{k=n}^{n+r-1} a_k \cos\left(k - \frac{r}{2}\right)x \right\}. \tag{3.1}$$

Proof. An elementary calculation gives

$$\sum_{k=n}^{2n-1} a_k \sin kx = \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \sin kx + \sum_{k=n}^{2n-1} a_{k+r} \sin kx$$

$$= \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \sin kx + \cos rx \sum_{k=n+r}^{2n+r-1} a_k \sin kx - \sin rx \sum_{k=n+r}^{2n+r-1} a_k \cos kx. \quad (3.2)$$

On the other hand

$$\sum_{k=n}^{2n-1} a_k \cos kx = \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \cos kx + \sum_{k=n}^{2n-1} a_{k+r} \cos kx$$

$$= \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \cos kx + \cos rx \sum_{k=n+r}^{2n+r-1} a_k \cos kx + \sin rx \sum_{k=n+r}^{2n+r-1} a_k \sin kx.$$

Hence

$$(1 - \cos rx) \sum_{k=n+r}^{2n+r-1} a_k \cos kx = \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \cos kx + \sin rx \sum_{k=n+r}^{2n+r-1} a_k \sin kx$$

$$- \sum_{k=n}^{n+r-1} a_k \cos kx + \sum_{k=2n}^{2n+r-1} a_k \cos kx.$$

Therefore, if $x \neq \frac{2l\pi}{r}$, then

$$\sum_{k=n+r}^{2n+r-1} a_k \cos kx = \frac{1}{2 \sin (rx/2)} \left\{ \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \cos kx \right.$$

$$\left. + \sin rx \sum_{k=n+r}^{2n+r-1} a_k \sin kx - \sum_{k=n}^{n+r-1} a_k \cos kx + \sum_{k=2n}^{2n+r-1} a_k \cos kx \right\}.$$

Putting this to (3.2) we get

$$\sum_{k=n}^{2n-1} a_k \sin kx = \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \sin kx + \cos rx \sum_{k=n+r}^{2n+r-1} a_k \sin kx$$

$$- \frac{\cos (rx/2)}{\sin (rx/2)} \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \cos kx - 2 \cos^2 \frac{rx}{2} \sum_{k=n+r}^{2n+r-1} a_k \sin kx$$

$$+ \frac{\cos (rx/2)}{\sin (rx/2)} \sum_{k=n}^{n+r-1} a_k \cos kx - \frac{\cos (rx/2)}{\sin (rx/2)} \sum_{k=2n}^{2n+r-1} a_k \cos kx$$

$$= \frac{-1}{\sin (rx/2)} \sum_{k=n}^{2n-1} (a_k - a_{k+r}) \cos \left(k + \frac{r}{2} \right) - \sum_{k=n+r}^{2n+r-1} a_k \sin kx$$

$$+ \frac{\cos (rx/2)}{\sin (rx/2)} \sum_{k=n}^{n+r-1} a_k \cos kx - \frac{\cos (rx/2)}{\sin (rx/2)} \sum_{k=2n}^{2n+r-1} a_k \cos kx.$$

Thus

$$2\sum_{k=n}^{2n-1} a_k \sin kx = \frac{-1}{\sin\left(rx/2\right)} \sum_{k=n}^{2n-1} \left(a_k - a_{k+r}\right) \cos\left(k + \frac{r}{2}\right)$$

$$+\frac{\cos\left(rx/2\right)}{\sin\left(rx/2\right)} \sum_{k=n}^{n+r-1} a_k \cos kx - \frac{\cos\left(rx/2\right)}{\sin\left(rx/2\right)} \sum_{k=2n}^{2n+r-1} a_k \cos kx$$

$$+\sum_{k=n}^{n+r-1} a_k \sin kx - \sum_{k=2n}^{2n+r-1} a_k \sin kx$$

$$= \frac{-1}{\sin\left(rx/2\right)} \left\{ \sum_{k=n}^{2n-1} \left(a_k - a_{k+r}\right) \cos\left(k + \frac{r}{2}\right) x \right.$$

$$\left. + \sum_{k=2n}^{2n+r-1} a_k \cos\left(k - \frac{r}{2}\right) x - \sum_{k=n}^{n+r-1} a_k \cos\left(k - \frac{r}{2}\right) x \right\}$$

and (3.1) holds.

The proof is complete. $\square$

Definition 3. A complex sequence $d := (d_n)$ is said to be weak monotone if

$$n\left|d_n\right| \leq K\left(d\right) \sum_{k=[n/c]}^{[cn]} \left|d_k\right|, \quad c > 1,$$

holds for all n.

Lemma 2. *[13, Theorem 2.2] Let a nonnegative sequence (b_n) be weak monotone. Then the uniform convergence of series (1.1) (or the fact that series (3.1) is the Fourier series of a continuous function) implies the condition $nb_n \to 0$ as $n \to \infty$.*

4 Proofs

In this section we shall prove our theorems and remarks.

4.1 Proof of Remark 1

Let $r \in \mathbb{N}$ and $(a_n) \in GM\left(\beta, 1\right)$. Then for all n

$$\sum_{k=n}^{2n-1} \left|a_k - a_{k+r}\right| = \sum_{k=n}^{2n-1} \left| \sum_{i=0}^{r-1} \left(a_{k+i} - a_{k+i+1}\right) \right|$$

$$\leq \sum_{k=n}^{2n-1} \sum_{i=0}^{r-1} |a_{k+i} - a_{k+i+1}| = \sum_{i=0}^{r-1} \sum_{k=n+i}^{2n+i-1} |a_k - a_{k+1}| \ll \sum_{i=0}^{r-1} \beta_{n+i} \ll \beta_n$$

and $(a_n) \in GM(\beta, r)$. $\Box$

4.2 Proof of Theorem 2

If $r_1 \mid r_2$, then exists a natural number p such that $r_2 = p \cdot r_1$. Supposing $(a_n) \in GM(\beta^*, r_1)$ we have for all n

$$\sum_{k=n}^{2n-1} |a_k - a_{k+r_2}| = \sum_{k=n}^{2n-1} \left| \sum_{l=0}^{p-1} a_{k+l\cdot r_1} - a_{k+(l+1)\cdot r_1} \right|$$

$$\leq \sum_{k=n}^{2n-1} \sum_{l=0}^{p-1} \left| a_{k+l\cdot r_1} - a_{k+(l+1)\cdot r_1} \right| = \sum_{l=0}^{p-1} \sum_{k=n+l\cdot r_1}^{2n+l\cdot r_1 - 1} |a_k - a_{k+r_1}|$$

$$\ll \sum_{l=0}^{p-1} \left(\sum_{k=n+l\cdot r_1}^{n+l\cdot r_1 + r_1 - 1} |a_k| + \sum_{k=[c^{-1}(n+l\cdot r_1)]}^{[c(n+l\cdot r_1)]} \frac{|a_k|}{k} \right)$$

$$\leq p \left(\sum_{k=n}^{n+p\cdot r_1 - 1} |a_k| + \sum_{k=[c^{-1}n]}^{[c(n+p\cdot r_1)]} \frac{|a_k|}{k} \right) \ll \sum_{k=n}^{n+r_2-1} |a_k| + \sum_{k=\left[(c(1+r_2))^{-1} n\right]}^{[c(1+r_2)n]} \frac{|a_k|}{k},$$

whence $(a_n) \in GM(\beta^*, r_2)$. Thus $GM(\beta^*, r_1) \subseteq GM(\beta^*, r_2)$.

Now, we prove that $GM(\beta^*, r_1) \neq GM(\beta^*, r_2)$.

Let

$$a_n = \frac{2 + \alpha_n}{n^2}, \quad \text{where} \quad \alpha_n = \begin{cases} -1 & \text{if} \quad r_2 \mid n, \\ 1 & \text{if} \quad r_2 \nmid n. \end{cases}$$

We show that $(a_n) \in GM(\beta^*, r_2)$ and $(a_n) \notin GM(\beta^*, r_1)$. Let

$$A_{r_2} := A(r_2, k, n) = \{k : n \leq k < 2n \quad \text{and} \quad r_2 \mid k\},$$

$$B_{r_2} := B(r_2, k, n) = \{k : n \leq k < 2n \quad \text{and} \quad r_2 \nmid k\}.$$

Then for all n

$$\sum_{k=n}^{2n-1} |a_k - a_{k+r_2}| = \left(\sum_{k \in A_{r_2}} + \sum_{k \in B_{r_2}} \right) |a_k - a_{k+r_2}|$$

$$= \sum_{k \in A_{r_2}} \left| \frac{1}{k^2} - \frac{1}{(k+r_2)^2} \right| + \sum_{k \in B_{r_2}} \left| \frac{3}{k^2} - \frac{3}{(k+r_2)^2} \right|$$

$$\leq 3\sum_{k=n}^{2n-1}\frac{2kr_2+r_2^2}{k^2\left(k+r_2\right)^2}\leq 6r_2\sum_{k=n}^{2n-1}\frac{1}{k^3}\leq\frac{6r_2}{n^2}\ll\sum_{k=[n/c]}^{[cn]}\frac{1}{k^3}\leq\sum_{k=[n/c]}^{[cn]}\frac{a_k}{k}$$

and we have $(a_n)\in GM\left(\beta^*,r_2\right).$

Since $r_1<r_2$ we obtain that $r_2\nmid r_1$ and

$$\sum_{k=n}^{2n-1}\left|a_k-a_{k+r_1}\right|\geq\sum_{k\in A_{r_2}}\left|a_k-a_{k+r_1}\right|=\sum_{k\in A_{r_2}}\left|\frac{1}{k^2}-\frac{3}{\left(k+r_1\right)^2}\right|$$

$$=\sum_{k\in A_{r_2}}\frac{\left|2k^2-2kr_1-r_1^2\right|}{k^2\left(k+r_1\right)^2}.$$

If $n\geq 5r_1$ then $2n^2-2nr_1-r_1^2\geq\left(n+r_1\right)^2$. Hence for $n\geq 5r_1$

$$\sum_{k=n}^{2n-1}\left|a_k-a_{k+r_1}\right|\geq\sum_{k\in A_{r_2}}\frac{1}{k^2}\geq\frac{1}{4nr_2}$$

and since

$$\sum_{k=n}^{n+r_1-1}a_k+\sum_{k=[n/c]}^{[cn]}\frac{a_k}{k}\leq\frac{3\left(r_1+c^4\right)}{n^2},$$

the inequality

$$\sum_{k=n}^{2n-1}\left|a_k-a_{k+r_1}\right|\leq K\left(a\right)\left\{\sum_{k=n}^{n+r_1-1}a_k+\sum_{k=[n/c]}^{[cn]}\frac{a_k}{k}\right\}$$

does not hold, that is, (a_n) does not belong to $GM\left(\beta^*,r_1\right).$

This complete the proof. □

4.3 Proof of Theorem 3

From the above proof we can see that there exists a sequence $(a_n)\in GM\left(\beta^*,r_2\right)$ such that $(a_n)\notin GM\left(\beta^*,r_1\right).$

Analogously, let

$$e_n=\frac{2+\gamma_n}{n^2},\quad\text{where}\quad\gamma_n=\begin{cases}-1 & \text{if}\quad r_1\mid n,\\ 1 & \text{if}\quad r_1\nmid n.\end{cases}$$

We show that $(e_n)\in GM\left(\beta^*,r_1\right)$ and $(e_n)\notin GM\left(\beta^*,r_2\right)$. Let

$$A_{r_1}:=A\left(r_1,k,n\right)=\left\{k:n\leq k<2n\quad\text{and}\quad r_1\mid k\right\},$$

$$B_{r_1} := B(r_1,k,n) = \{k : n \le k < 2n \quad \text{and} \quad r_1 \nmid k\}.$$

Then for all n

$$\sum_{k=n}^{2n-1} |e_k - e_{k+r_1}| = \left(\sum_{k\in A_{r_1}} + \sum_{k\in B_{r_1}} \right) |e_k - e_{k+r_1}|$$

$$= \sum_{k\in A_{r_1}} \left| \frac{1}{k^2} - \frac{1}{(k+r_1)^2} \right| + \sum_{k\in B_{r_1}} \left| \frac{3}{k^2} - \frac{3}{(k+r_1)^2} \right|$$

$$\le 3 \sum_{k=n}^{2n-1} \frac{2kr_1 + r_1^2}{k^2 (k+r_1)^2} \le 6r_1 \sum_{k=n}^{2n-1} \frac{1}{k^3} \le \frac{6r_1}{n^2} \ll \sum_{k=[n/c]}^{[cn]} \frac{1}{k^3} \le \sum_{k=[n/c]}^{[cn]} \frac{e_k}{k}$$

and $(e_n) \in GM(\beta^*, r_1)$.

If $r_1 \nmid r_2$ then for $n \ge 5r_2$ we get

$$\sum_{k=n}^{2n-1} |e_k - e_{k+r_2}| \ge \sum_{k\in A_{r_1}} |e_k - e_{k+r_2}| = \sum_{k\in A_{r_1}} \left| \frac{1}{k^2} - \frac{3}{(k+r_1)^2} \right|$$

$$= \sum_{k\in A_{r_1}} \frac{\left|2k^2 - 2kr_2 - r_2^2\right|}{k^2 (k+r_2)^2} \ge \sum_{k\in A_{r_1}} \frac{1}{k^2} \ge \frac{1}{4nr_1}$$

and since

$$\sum_{k=n}^{n+r_2-1} e_k + \sum_{k=[n/c]}^{[cn]} \frac{e_k}{k} \le \frac{3\left(r_2 + c^4\right)}{n^2},$$

the inequality

$$\sum_{k=n}^{2n-1} |e_k - e_{k+r_1}| \le K(e) \left\{ \sum_{k=n}^{n+r_2-1} e_k + \sum_{k=[n/c]}^{[cn]} \frac{e_k}{k} \right\}$$

does not hold, that is, (e_n) does not belong to $GM(\beta^*, r_2)$.

This complete the proof. □

4.4 Proof of Theorem 4

Let $(b_n) \in GM(\beta^*, r)$, where $r \ge 1$. We show that such sequence (b_n) is weak monotone. If $n \le r$ then the inequality obviously holds.

Now, let $n > r$. For $j = n+1, n+2, ..., 2n$ we get

$$\sum_{k=n}^{j-1} |b_k - b_{k+r}| \ge \sum_{k=n}^{j-1} b_k - \sum_{k=n+r}^{j+r-1} b_k$$

and for $j \geq n+r+1$ we obtain

$$b_n \leq \sum_{k=n}^{n+r-1} b_k \leq \sum_{k=n}^{j-1} |b_k - b_{k+r}| + \sum_{k=j}^{j+r-1} b_k$$

$$\leq \sum_{k=[j/2]}^{2[j/2]-1} |b_k - b_{k+r}| + \sum_{k=j}^{j+r-1} b_k$$

$$\ll \sum_{k=[j/2c]}^{[cj/2]} \frac{b_k}{k} + \sum_{k=[j/2]}^{[j/2]+r-1} b_k + \sum_{k=j}^{j+r-1} b_k \leq \frac{1}{n} \sum_{k=[n/2c]}^{[cn]} b_k + \sum_{k=[j/2]}^{[j/2]+r-1} b_k + \sum_{k=j}^{j+r-1} b_k.$$

Summing up on j we get

$$nb_n = \sum_{j=n+1}^{2n} b_n = \sum_{j=n+1}^{n+r} b_n + \sum_{j=n+r+1}^{2n} b_n$$

$$\ll rb_n + \sum_{j=n+r+1}^{2n} \left(\frac{1}{n} \sum_{k=[n/2c]}^{[cn]} b_k + \sum_{k=[j/2]}^{[j/2]+r-1} b_k + \sum_{k=j}^{j+r-1} b_k \right)$$

$$\leq rb_n + \sum_{k=[n/2c]}^{[cn]} b_k + \sum_{j=n+1}^{2n} \sum_{k=j}^{j+r-1} b_k + \sum_{j=n+1}^{2n} \sum_{k=[j/2]}^{[j/2]+r-1} b_k$$

$$\leq r \sum_{k=[n/2c]}^{[cn]} b_k + \sum_{j=0}^{r-1} \sum_{k=n+1+j}^{2n-1+j} b_k + 2 \sum_{j=0}^{r-1} \sum_{k=[(n+1)/2]+j}^{[(2n-1)/2]+j} b_k$$

$$\leq r \sum_{k=[n/2c]}^{[cn]} b_k + r \sum_{k=n+1}^{2n+r-2} b_k + 2r \sum_{k=[(n+1)/2]}^{[(2n-1)/2]+r-1} b_k$$

$$\leq r \sum_{k=[n/2c]}^{[cn]} b_k + 3r \sum_{k=[n/2]}^{2n+r} b_k \leq r \sum_{k=[n/2c]}^{[2cn]} b_k + 3r \sum_{k=[n/2c]}^{3n} b_k \leq 4r \sum_{k=[n/c_1]}^{[nc_1]} b_k,$$

where $c_1 = \max\{3, 2c\}$. Therefore (b_n) is weak monotone and by Lemma 2 we have that $nb_n \to 0$ as $n \to \infty$ and thus the proof is complete. $\square$

4.5 Proof of Theorem 5

Denote

$$\varepsilon_n^{(1)} := \sup_{k \geq n/c} k\,|b_k|, \quad \text{where} \quad c > 1$$

and

$$r_n(x)=\sum_{k=n}^{\infty}b_k\sin kx.$$

In view of the assumption, we can see that $\varepsilon_n^{(1)}\to 0$ as $n\to\infty$. Further, we shall show that

$$|r_n(x)|\ll\varepsilon_n^{(1)}. \tag{4.1}$$

Since $r_n(l\pi)=0$, where $l\in\mathbb{Z}$, it suffices to prove (4.1) for $0<x<\pi$.

First we show that (4.1) is valid for $0<x\leq\frac{\pi}{2}$.

Let $N:=N(x)\geq 2$ be the natural number such that

$$\frac{\pi}{N+1}<x\leq\frac{\pi}{N}. \tag{4.2}$$

Then

$$r_n(x)=\sum_{k=n}^{n+N-1}b_k\sin kx+\sum_{k=n+N}^{\infty}b_k\sin kx=r_n^{(1)}(x)+r_n^{(2)}(x).$$

Hence, by (4.2),

$$\left|r_n^{(1)}(x)\right|\leq\sum_{k=n}^{n+N-1}|b_k|\,|\sin kx|\leq x\sum_{k=n}^{n+N-1}k\,|b_k|\leq x\varepsilon_n^{(1)}N\leq\pi\varepsilon_n^{(1)}. \tag{4.3}$$

If $(b_n)\in GM(\beta^*,2)$, then using Lemma 1, the inequality $\frac{2}{\pi}x\leq\sin x$ $\left(x\in\left[0,\frac{\pi}{2}\right]\right)$ and (4.2) we obtain

$$\left|r_n^{(2)}(x)\right|=\left|\sum_{j=0}^{\infty}\sum_{k=2^j(n+N)}^{2^{j+1}(n+N)-1}b_k\sin kx\right|$$

$$\leq\sum_{j=0}^{\infty}\left|\frac{-1}{2\sin x}\sum_{k=2^j(n+N)}^{2^{j+1}(n+N)-1}(b_k-b_{k+2})\cos(k+1)x\right.$$

$$\left.+\sum_{k=2^{j+1}(n+N)}^{2^{j+1}(n+N)+1}b_k\cos(k-1)x-\sum_{k=2^j(n+N)}^{2^j(n+N)+1}b_k\cos(k-1)x\right\}\Bigg|$$

$$\leq\frac{1}{2\sin x}\sum_{j=0}^{\infty}\left\{\sum_{k=2^j(n+N)}^{2^{j+1}(n+N)-1}|b_k-b_{k+2}|+\sum_{k=2^{j+1}(n+N)}^{2^{j+1}(n+N)+1}|b_k|+\sum_{k=2^j(n+N)}^{2^j(n+N)+1}|b_k|\right\}$$

$$\ll\frac{\pi}{2x}\sum_{j=0}^{\infty}\left\{\sum_{k=2^j(n+N)}^{2^{j+1}(n+N)-1}|b_k-b_{k+2}|+\sum_{k=2^j(n+N)}^{2^j(n+N)+1}|b_k|\right\}$$

$$\ll (N+1)\sum_{j=0}^{\infty}\left\{\sum_{k=2^j(n+N)}^{2^j(n+N)+1}|b_k|+\sum_{k=[2^j(n+N)/c]}^{\left[c2^j(n+N)\right]}\frac{|b_k|}{k}\right\}$$

$$\ll (N+1)\,\varepsilon_n^{(1)}\sum_{j=0}^{\infty}\left\{\sum_{k=2^j(n+N)}^{2^j(n+N)+1}\frac{1}{k}+\sum_{k=[2^j(n+N)/c]}^{\left[c2^j(n+N)\right]}\frac{1}{k^2}\right\}$$

$$\ll \frac{N+1}{N+n}\varepsilon_n^{(1)}\sum_{j=0}^{\infty}\frac{1}{2^j}\ll\varepsilon_n^{(1)}. \tag{4.4}$$

Now, we prove (4.1) for $\frac{\pi}{2}\le x<\pi$.

Let $M:=M(x)\ge 2$ be the natural number such that

$$\pi-\frac{\pi}{M}\le x<\pi-\frac{\pi}{M+1}. \tag{4.5}$$

Then

$$r_n(x)=\sum_{k=n}^{n+M-1}b_k\sin kx+\sum_{k=n+M}^{\infty}b_k\sin kx=r_n^{(3)}(x)+r_n^{(4)}(x).$$

Using the inequality $\sin x\le\pi-x$ $(x\in(0,\pi))$ and (4.5) we get

$$\left|r_n^{(3)}(x)\right|\le(\pi-x)\sum_{k=n}^{n+M-1}k\,|b_k|\le(\pi-x)\,M\varepsilon_n^{(1)}\le\pi\varepsilon_n^{(1)}. \tag{4.6}$$

If $(b_n)\in GM(\beta^*,2)$, then using Lemma 1, the inequality $2-\frac{2}{\pi}x\le\sin x$ $\left(x\in\left[\frac{\pi}{2},\pi\right]\right)$ and (4.5) we obtain

$$\left|r_n^{(2)}(x)\right|=\left|\sum_{j=0}^{\infty}\sum_{k=2^j(n+M)}^{2^{j+1}(n+M)-1}b_k\sin kx\right|$$

$$\le\sum_{j=0}^{\infty}\left|\frac{-1}{2\sin x}\sum_{k=2^j(n+M)}^{2^{j+1}(n+M)-1}(b_k-b_{k+2})\cos(k+1)x\right.$$

$$\left.+\sum_{k=2^{j+1}(n+M)}^{2^{j+1}(n+M)+1}b_k\cos(k-1)x-\sum_{k=2^j(n+M)}^{2^j(n+M)+1}b_k\cos(k-1)x\right\}\Bigg|$$

$$\ll\frac{1}{2\left(1-\frac{1}{\pi}x\right)}\sum_{j=0}^{\infty}\left\{\sum_{k=2^j(n+M)}^{2^{j+1}(n+M)-1}|b_k-b_{k+2}|+\sum_{k=2^j(n+M)}^{2^j(n+M)+1}|b_k|\right\}$$

$$\ll(M+1)\sum_{j=0}^{\infty}\left\{\sum_{k=2^j(n+M)}^{2^j(n+M)+1}|b_k|+\sum_{k=[2^j(n+M)/c]}^{\left[c2^j(n+M)\right]}\frac{|b_k|}{k}\right\}$$

$$\ll \frac{M+1}{M+n}\varepsilon_n^{(1)}\sum_{j=0}^{\infty}\frac{1}{2^j} \ll \varepsilon_n^{(1)}. \tag{4.7}$$

From the estimations (4.3), (4.4), (4.6) and (4.7)we obtain the uniform convergence of series (1.1) follows and thus the proof is complete. □

4.6 Proof of Remark 3

Let $x_0 = \frac{2\pi}{3}$ and

$$d_n = \begin{cases} \frac{3}{n\ln(n+1)} & \text{if} \quad n = 3k+1, \\ \frac{1}{n\ln(n+1)} & \text{if} \quad n \neq 3k+1, \end{cases}$$

where $k \in \mathbb{N}\cup\{0\}$. An elementary calculation gives

$$\sum_{k=1}^{\infty} d_k \sin(kx_0) = d_1 \sin\frac{2\pi}{2} + d_2\sin\frac{4\pi}{2} + \sum_{k=1}^{\infty}\sum_{i=0}^{2} d_{3k+i}\sin(3k+i)\frac{2\pi}{3}$$

$$= (d_1 - d_2)\sin\frac{2\pi}{3} + \sum_{k=1}^{\infty}\sum_{i=0}^{2} d_{3k+i}\sin\left(i\frac{2\pi}{3}\right)$$

$$= (d_1 - d_2)\sin\frac{2\pi}{3} + \sum_{k=1}^{\infty}\sum_{i=1}^{2} d_{3k+i}\sin\left(i\frac{2\pi}{3}\right) = \sin\frac{2\pi}{3}\sum_{k=1}^{\infty}(d_{3k+1} - d_{3k+2})$$

$$= \sin\frac{2\pi}{3}\sum_{k=1}^{\infty}\left(\frac{3}{(3k+1)\ln(3k+2)} - \frac{1}{(3k+2)\ln(3k+3)}\right)$$

$$\geq \sin\frac{2\pi}{3}\sum_{k=1}^{\infty}\frac{2}{(3k+1)\ln(3k+3)} = \infty$$

and the series $\sum_{k=1}^{\infty} d_k\sin(kx_0)$ is divergent, too. □

4.7 Proof of Theorem 6

Denote

$$\varepsilon_n^{(1)} := \sup_{k\geq n/c} k\,|b_k|\,, \quad \text{where} \quad c>1,$$

$$\varepsilon_n^{(2)} := \sum_{k\geq n/r}^{\infty}\sum_{k=1}^{[r/2]}|b_{r\cdot n+k} - b_{r\cdot n+r-k}|$$

and

$$r_n(x) = \sum_{k=n}^{\infty} b_k \sin kx.$$

In view of the assumptions, we have that $\varepsilon_n^{(1)} \to 0$ and $\varepsilon_n^{(2)} \to 0$ as $n \to \infty$. We shall show that

$$|r_n(x)| \ll \varepsilon_n^{(1)} + \varepsilon_n^{(2)} \tag{4.8}$$

also holds. Since $r_n(l\pi) = 0$, where $l \in \mathbb{Z}$, it suffices to prove (4.8) for $0 < x < \pi$.

First we shall that (4.8) is valid for $x = \frac{2l\pi}{r}$, where l is an integer number such that $0 < 2l < r$. For any n exist two number $p, q \in \mathbb{N} \cup \{0\}$ such that $n = r \cdot p + q$, where $0 \leq q < r$. Then

$$r_n\left(\frac{2l\pi}{r}\right) = \sum_{k=n}^{\infty} b_k \sin\left(k\frac{2l\pi}{r}\right) = \sum_{k=r\cdot p+q}^{r(p+1)-1} b_k \sin\left(k\frac{2l\pi}{r}\right)$$

$$+ \sum_{k=p+1}^{\infty} \sum_{i=0}^{r-1} b_{r\cdot k+i} \sin\left((rk+i)\frac{2l\pi}{r}\right) = \sum_{k=r\cdot p+q}^{r(p+1)-1} b_k \sin\left(k\frac{2l\pi}{r}\right)$$

$$+ \sum_{k=p+1}^{\infty} \sum_{i=1}^{r-1} b_{r\cdot k+i} \sin\left(i\frac{2l\pi}{r}\right).$$

If $r = 2s$ $(s = 2, 3, ..)$ then

$$\sum_{i=1}^{r-1} b_{r\cdot k+i} \sin\left(i\frac{2l\pi}{r}\right) = \sum_{k=1}^{2s-1} b_{2s\cdot k+i} \sin\left(i\frac{l\pi}{s}\right) = \sum_{i=1}^{s-1} \left(b_{2s\cdot k+i} - b_{2s\cdot k+2s-i}\right) \sin\left(i\frac{l\pi}{s}\right)$$

$$= \sum_{i=1}^{s} \left(b_{2s\cdot k+i} - b_{2s\cdot k+2s-i}\right) \sin\left(i\frac{l\pi}{s}\right) = \sum_{i=1}^{r/2} \left(b_{r\cdot k+i} - b_{r\cdot k+r-i}\right) \left(i\frac{2l\pi}{r}\right) \tag{4.9}$$

and if $r = 2s+1$ $(s = 1, 2, ..)$ then

$$\sum_{i=1}^{r-1} b_{r\cdot k+i} \sin\left(i\frac{2l\pi}{r}\right) = \sum_{k=1}^{2s} b_{(2s+1)\cdot k+i} \sin\left(i\frac{2l\pi}{2s+1}\right)$$

$$= \sum_{k=1}^{s} \left(b_{(2s+1)\cdot k+i} - b_{(2s+1)\cdot k+2s+1-i}\right) \left(i\frac{2l\pi}{2s+1}\right)$$

$$= \sum_{i=1}^{[r/2]} \left(b_{r\cdot k+i} - b_{r\cdot k+r-i}\right) \left(i\frac{2l\pi}{r}\right). \tag{4.10}$$

Using (4.9) or (4.10) we obtain

$$\left|r_n\left(\frac{2l\pi}{r}\right)\right| \leq \sum_{k=r\cdot p+q}^{r(p+1)-1} |b_k| + \sum_{k=p+1}^{\infty} \sum_{i=1}^{[r/2]} |b_{r\cdot k+i} - b_{r\cdot k+r-i}|$$

$$\leq \frac{r}{n}\varepsilon_n^{(1)} + \varepsilon_n^{(2)} \ll \varepsilon_n^{(1)} + \varepsilon_n^{(2)}. \tag{4.11}$$

Now, we prove that (4.8) holds for $\frac{2l\pi}{r} < x \leq \frac{2l\pi}{r} + \frac{\pi}{r}$, where $0 \leq 2l < r$.

Let $N := N(x)$ be the natural number such that

$$\frac{2l\pi}{r} + \frac{\pi}{N+1} < x \leq \frac{2l\pi}{r} + \frac{\pi}{N}. \tag{4.12}$$

Then

$$r_n(x) = \sum_{k=n}^{n+N-1} b_k \sin kx + \sum_{k=n+N}^{\infty} b_k \sin kx = r_n^{(1)}(x) + r_n^{(2)}(x).$$

Applying Lagrange's mean value theorem to the function $f(x) = \sin kx$ on the interval $\left[\frac{2l\pi}{r}, x\right]$ we obtain that there exists $y \in \left(\frac{2l\pi}{r}, x\right)$ such that

$$\sin kx - \sin\left(k\frac{2l\pi}{r}\right) = k\cos ky\left(x - \frac{2l\pi}{r}\right).$$

Using this we get

$$\begin{aligned} r_n^{(1)}(x) &= \sum_{k=n}^{n+N-1} kb_k \cos ky\left(x - \frac{2l\pi}{r}\right) + \sum_{k=n}^{n+N-1} b_k \sin\left(k\frac{2l\pi}{r}\right) \\ &= r_n^{(1.1)}(x) + r_n^{(1.2)}\left(\frac{2l\pi}{r}\right). \end{aligned}$$

Hence, by (4.12),

$$\left|r_n^{(1.1)}(x)\right| \leq \left(x - \frac{2l\pi}{r}\right) \sum_{k=n}^{n+N-1} k|b_k| \leq \pi\varepsilon_n^{(1)}. \tag{4.13}$$

If $l = 0$ then $r_n^{(1.2)}(0) = 0$ and (4.8) is evident. Let $0 < 2l < r$. For any n and $N \geq r$ there exist four numbers $p_1, p_2, q_1, q_2 \in \mathbb{N} \cup \{0\}$ such that $n = rp_1 + q_1$ and $N = rp_2 + q_2$, where $p_2 \geq 1$ and $0 \leq q_1, q_2 < r$. If $q_1 + q_2 = r$ then

$$\begin{aligned} r_n^{(1.2)}\left(\frac{2l\pi}{r}\right) &= \sum_{k=r\cdot p_{1+q_1}}^{r(p_1+1)-1} b_k \sin\left(k\frac{2l\pi}{r}\right) + \sum_{k=r(p_1+1)}^{r(p_1+p_2)+r-1} b_k \sin\left(k\frac{2l\pi}{r}\right) \\ &= \sum_{k=r\cdot p_{1+q_1}}^{r(p_1+1)-1} b_k \sin\left(k\frac{2l\pi}{r}\right) + \sum_{k=p_1+1}^{p_1+p_2} \sum_{i=0}^{r-1} b_{r\cdot k+i} \sin\left(i\frac{2l\pi}{r}\right) \end{aligned} \tag{4.14}$$

and using (4.9) or (4.10) we obtain

$$\left|r_n^{(1.2)}\left(\frac{2l\pi}{r}\right)\right| \leq \sum_{k=r\cdot p_1+q_1}^{r(p_1+1)-1} |b_k| + \sum_{k=p_1+1}^{p_1+p_2} \sum_{i=1}^{[r/2]} |b_{r\cdot k+i} - b_{r\cdot k+r-i}| \ll \varepsilon_n^{(1)} + \varepsilon_n^{(2)}.$$

Let $q_1 + q_2 < r$. Then

$$r_n^{(1.2)}\left(\frac{2l\pi}{r}\right) = \sum_{k=r\cdot p_1+q_1}^{r(p_1+1)-1} b_k \sin\left(k\frac{2l\pi}{r}\right)$$

$$+\sum_{k=p_1+1}^{p_1+p_2}\sum_{i=0}^{r-1} b_{r\cdot k+i}\sin\left(i\frac{2l\pi}{r}\right) - \sum_{k=r(p_1+p_2)+q_1+q_2-1}^{r(p_1+p_2)+r-1} b_k \sin\left(k\frac{2l\pi}{r}\right)$$

and by (4.9) or (4.10)

$$\left|r_n^{(1.2)}\left(\frac{2l\pi}{r}\right)\right| \le \sum_{k=r\cdot p+q_1}^{r(p_1+1)-1} |b_k| + \sum_{k=p_1+1}^{p_1+p_2}\sum_{i=1}^{[r/2]} |b_{r\cdot k+i} - b_{r\cdot k+r-i}|$$

$$+\sum_{k=r(p_1+p_2)+q_1+q_2-1}^{r(p_1+p_2)+r-1} |b_k| \ll \varepsilon_n^{(1)} + \varepsilon_n^{(2)}. \tag{4.15}$$

Now, assume $q_1 + q_2 > r$. Then

$$r_n^{(1.2)}\left(\frac{2l\pi}{r}\right) = \sum_{k=r\cdot p_{1+q_1}}^{r(p_1+1)-1} b_k \sin\left(k\frac{2l\pi}{r}\right)$$

$$+\sum_{k=p_1+1}^{p_1+p_2}\sum_{i=0}^{r-1} b_{r\cdot k+i}\sin\left(i\frac{2l\pi}{r}\right) + \sum_{k=r(p_1+p_2)+r}^{r(p_1+p_2)+q_1+q_2-1} b_k \sin\left(k\frac{2l\pi}{r}\right)$$

and using (4.9) or (4.10) we have

$$\left|r_n^{(1.2)}\left(\frac{2l\pi}{r}\right)\right| \le \sum_{k=r\cdot p_1+q_1}^{r(p_1+1)-1} |b_k| + \sum_{k=p_1+1}^{p_1+p_2}\sum_{i=1}^{[r/2]} |b_{r\cdot k+i} - b_{r\cdot k+r-i}|$$

$$+\sum_{k=r(p_1+p_2)+r}^{r(p_1+p_2)+q_1+q_2-1} |b_k| \ll \varepsilon_n^{(1)} + \varepsilon_n^{(2)}. \tag{4.16}$$

Therefore, by (4.14)-(4.15),

$$\left|r_n^{(1.2)}\left(\frac{2l\pi}{r}\right)\right| \ll \varepsilon_n^{(1)} + \varepsilon_n^{(2)}. \tag{4.17}$$

If $(b_n) \in GM(\beta^*, 3)$ $(r \ge 3)$, then using Lemma 1, the inequality $\frac{r}{\pi}x - 2l \le \left|\sin\frac{rx}{2}\right|$ $\left(x \in \left[\frac{2l\pi}{r}, \frac{2l\pi}{r} + \frac{\pi}{r}\right] \text{ and } 0 \le 2l < r\right)$ and (4.12) we obtain

$$\left|r_n^{(2)}(x)\right| = \left|\sum_{j=0}^{\infty}\sum_{k=2^j(n+N)}^{2^{j+1}(n+N)-1} b_k \sin kx\right|$$

$$\le \sum_{j=0}^{\infty}\Bigg|\frac{-1}{2\sin\left(rx/2\right)}\sum_{k=2^j(n+N)}^{2^{j+1}(n+N)-1}\left(b_k-b_{k+r}\right)\cos\left(k+\frac{r}{2}\right)x$$

$$+\sum_{k=2^{j+1}(n+N)}^{2^{j+1}(n+N)+r-1} b_k\cos\left(k-\frac{r}{2}\right)x-\sum_{k=2^j(n+N)}^{2^j(n+N)+r-1} b_k\cos\left(k-\frac{r}{2}\right)x\Bigg\}\Bigg|$$

$$\le \frac{1}{2\left|\sin\left(rx/2\right)\right|}\sum_{j=0}^{\infty}\left\{\sum_{k=2^j(n+N)}^{2^{j+1}(n+N)-1}\left|b_k-b_{k+r}\right|+\sum_{k=2^{j+1}(n+N)}^{2^{j+1}(n+N)+r-1}\left|b_k\right|+\sum_{k=2^j(n+N)}^{2^j(n+N)+r-1}\left|b_k\right|\right\}$$

$$\ll \frac{1}{\frac{r}{\pi}x-2l}\sum_{j=0}^{\infty}\left\{\sum_{k=2^j(n+N)}^{2^{j+1}(n+N)-1}\left|b_k-b_{k+r}\right|+\sum_{k=2^j(n+N)}^{2^j(n+N)+r-1}\left|b_k\right|\right\}$$

$$\ll \frac{N+1}{r}\sum_{j=0}^{\infty}\left\{\sum_{k=2^j(n+N)}^{2^j(n+N)+r-1}\left|b_k\right|+\sum_{k=\left[2^j(n+N)/c\right]}^{\left[c2^j(n+N)\right]}\frac{\left|b_k\right|}{k}\right\}$$

$$\ll \frac{N+1}{N+n}\varepsilon_n^{(1)}\sum_{j=0}^{\infty}\frac{1}{2^j}\ll \varepsilon_n^{(1)}. \qquad (4.18)$$

Finally, we prove that (4.8) is true for $\frac{2l\pi}{r}+\frac{\pi}{r}\le x<\frac{2(l+1)\pi}{r}$, where $0<2\left(l+1\right)\le r$.

Let $M:=M\left(x\right)\ge r$ be the natural number such that

$$\frac{2\left(l+1\right)\pi}{r}-\frac{\pi}{M}\le x<\frac{2\left(l+1\right)\pi}{r}-\frac{\pi}{M+1}. \qquad (4.19)$$

Then

$$r_n\left(x\right)=\sum_{k=n}^{n+M-1} b_k\sin kx+\sum_{k=n+M}^{\infty} b_k\sin kx=r_n^{(3)}\left(x\right)+r_n^{(4)}\left(x\right).$$

Applying Lagrange's mean value theorem to the function $f\left(x\right)=\sin kx$ on the interval $\left[x,\frac{2(l+1)\pi}{r}\right]$ we obtain that there exists $z\in\left(x,\frac{2(l+1)\pi}{r}\right)$ such that

$$\sin\left(k\frac{2\left(l+1\right)\pi}{r}\right)-\sin kx=k\cos kz\left(\frac{2\left(l+1\right)\pi}{r}-x\right).$$

Using this we get

$$r_n^{(3)}\left(x\right)=\sum_{k=n}^{n+M-1} kb_k\cos kz\left(x-\frac{2\left(l+1\right)\pi}{r}\right)+\sum_{k=n}^{n+M-1} b_k\sin\left(k\frac{2\left(l+1\right)\pi}{r}\right)$$

$$=r_n^{(3.1)}\left(x\right)+r_n^{(3.2)}\left(\frac{2\left(l+1\right)\pi}{r}\right).$$

Hence, by (4.19),

$$\left|r_n^{(3.1)}(x)\right| \le \left(\frac{2(l+1)\pi}{r} - x\right) \sum_{k=n}^{n+M-1} k\,|b_k| \le \pi\varepsilon_n^{(1)}. \tag{4.20}$$

The quantity $r_n^{(3.2)}\left(\frac{2(l+1)\pi}{r}\right)$ we can estimate in the same way as the quantity $r_n^{(1.2)}\left(\frac{2l\pi}{r}\right)$. Therefore we get

$$\left|r_n^{(3.2)}\left(\frac{2(l+1)\pi}{r}\right)\right| \ll \varepsilon_n^{(1)} + \varepsilon_n^{(2)}. \tag{4.21}$$

If $(b_n) \in GM(\beta^*, 3)$ $(r \ge 3)$, then using Lemma 1, the inequality $2(l+1) - \frac{r}{\pi}x \le \left|\sin\frac{rx}{2}\right|$ $\left(x \in \left[\frac{2l\pi}{r} + \frac{\pi}{r}, \frac{2(l+1)\pi}{r}\right] \text{ and } 0 < 2(l+1) \le r\right)$ and (4.19) we obtain

$$\left|r_n^{(4)}(x)\right| = \left|\sum_{j=0}^{\infty} \sum_{k=2^j(n+M)}^{2^{j+1}(n+M)-1} b_k \sin kx\right|$$

$$\le \sum_{j=0}^{\infty} \left|\frac{-1}{2\sin(rx/2)} \sum_{k=2^j(n+M)}^{2^{j+1}(n+M)-1} (b_k - b_{k+r}) \cos\left(k + \frac{r}{2}\right)x\right.$$

$$\left. + \sum_{k=2^{j+1}(n+M)}^{2^{j+1}(n+M)+r-1} b_k \cos\left(k - \frac{r}{2}\right)x - \sum_{k=2^j(n+M)}^{2^j(n+M)+r-1} b_k \cos\left(k - \frac{r}{2}\right)x \right\}\right|$$

$$\le \frac{1}{2|\sin(rx/2)|} \sum_{j=0}^{\infty} \left\{ \sum_{k=2^j(n+M)}^{2^{j+1}(n+M)-1} |b_k - b_{k+r}| + \sum_{k=2^{j+1}(n+M)}^{2^{j+1}(n+M)+r-1} |b_k| + \sum_{k=2^j(n+M)}^{2^j(n+M)+r-1} |b_k| \right\}$$

$$\ll \frac{1}{2(l+1) - \frac{r}{\pi}x} \sum_{j=0}^{\infty} \left\{ \sum_{k=2^j(n+M)}^{2^{j+1}(n+M)-1} |b_k - b_{k+r}| + \sum_{k=2^j(n+M)}^{2^j(n+M)+r-1} b_k \right\}$$

$$\ll \frac{M+1}{r} \sum_{j=0}^{\infty} \left\{ \sum_{k=2^j(n+M)}^{2^j(n+M)+r-1} |b_k| + \sum_{k=[2^j(n+M)/c]}^{[c2^j(n+M)]} \frac{|b_k|}{k} \right\}$$

$$\ll \frac{M+1}{M+n} \varepsilon_n^{(1)} \sum_{j=0}^{\infty} \frac{1}{2^j} \ll \varepsilon_n^{(1)}. \tag{4.22}$$

From the estimation (4.11), (4.13), (4.17), (4.18), (4.20), (4.21) and (4.22) we obtain the uniform convergence of series (1.1) follows and thus the proof is complete. □

4.8 Proof of Remark 4

Let $r \geq 3$ and

$$a_n = \begin{cases} 0 & \text{if} \quad r \mid n, \\ \frac{1}{n \ln(n+1)} & \text{if} \quad r \nmid n. \end{cases}$$

It is clear that $na_n \to 0$ as $n \to \infty$.

First, we prove that $(a_n) \in GM\left(\beta^*, r\right)$. Let

$$A_r := A\left(r, k, n\right) = \left\{k : n \leq k < 2n \text{ and } r \mid k\right\}$$

and

$$B_r := B\left(r, k, n\right) \left\{k : n \leq k < 2n \text{ and } r \nmid k\right\}.$$

Then for all n

$$\sum_{k=n}^{2n-1} \left|a_k - a_{k+r}\right| = \left(\sum_{k \in A_r} + \sum_{k \in B_r}\right) \left|a_k - a_{k+r}\right|$$

$$= \sum_{k \in B_r} \left|\frac{1}{k \ln\left(k+1\right)} - \frac{1}{\left(k+r\right) \ln\left(k+r+1\right)}\right|$$

$$= \sum_{k \in B_r} \frac{k \ln\left(1 + \frac{r}{k+1}\right) + r \ln\left(k+r+1\right)}{k\left(k+r\right) \ln\left(k+1\right) \ln\left(k+r+1\right)} \leq 2r \sum_{k \in B_r} \frac{1}{k^2 \ln\left(k+1\right)}$$

$$= 2r \sum_{k=n}^{2n-1} \frac{a_k}{k} \ll \sum_{k=[n/c]}^{[cn]} \frac{a_k}{k}$$

and $(a_n) \in GM\left(\beta^*, r\right)$.

Now, we show that (2.1) is valid. We have

$$\sum_{n=1}^{\infty} \sum_{k=1}^{[r/2]} \left|a_{r \cdot n+k} - a_{r \cdot n+r-k}\right|$$

$$= \sum_{n=1}^{\infty} \sum_{k=1}^{[r/2]} \left|\frac{1}{\left(rn+k\right) \ln\left(rn+k+1\right)} - \frac{1}{\left(rn+r-k\right) \ln\left(rn+r-k+1\right)}\right|$$

$$\leq \sum_{n=1}^{\infty} \sum_{k=1}^{[r/2]} \frac{rn \ln\left(1 + \frac{r-2k}{rn+k+1}\right) + r \ln\left(rn+r-k+1\right)}{\left(rn+k\right)\left(rn+r-k\right) \ln\left(rn+k+1\right) \ln\left(rn+r-k+1\right)}$$

$$\leq \sum_{n=1}^{\infty} \sum_{k=1}^{[r/2]} \frac{r-2k + r \ln\left(rn+r-k+1\right)}{\left(rn+k\right)\left(rn+r-k\right) \ln\left(rn+k+1\right) \ln\left(rn+r-k+1\right)}$$

$$\leq 2r\sum_{n=1}^{\infty}\sum_{k=1}^{[r/2]}\frac{1}{(rn+k)\,(rn+r-k)\ln(rn+k+1)}\leq\sum_{n=1}^{\infty}\frac{1}{n^2\ln(n+1)}$$

and since the series $\sum\limits_{n=1}^{\infty}\frac{1}{n^2\ln(n+1)}$ converges (2.1) holds.

Finally, we prove that $(a_n)\notin GM(\beta^*,2)$. For all n we get

$$\sum_{k=n}^{2n-1}|a_k-a_{k+2}|\geq\sum_{k\in A_r}|a_k-a_{k+2}|=\sum_{k\in A_r}\frac{1}{(k+2)\ln(k+3)}$$
$$\geq\frac{1}{3r\ln(2n+2)}$$

and since

$$\sum_{k=n}^{n+1}a_k+\sum_{k=[n/c]}^{[cn]}\frac{a_k}{k}\ll\frac{1}{n\ln(n+1)},$$

the inequality

$$\sum_{k=n}^{2n-1}|a_k-a_{k+2}|\leq K(a)\left\{\sum_{k=n}^{n+1}a_k+\sum_{k=[n/c]}^{[cn]}\frac{a_k}{k}\right\}$$

does not hold, that is, (a_n) does not belong to $GM(\beta^*,2)$.

This ends our proof. □

References

[1] T. W. Chaundy and A. E. Jolliffe, The uniform convergence of a certain class of trigonometrical series, Proc. London Math. Soc., 15(1916), 214-216.

[2] R. J. Le and S. P. Zhou, A new condition for the uniform convergence of certain trigonometric series, Acta Math. Hungar., 108 (1-2) (2005), 161-169.

[3] L Leindler, On the uniform convergence and boundedness of a certain class of sine series, Analysis Math., 27 (2001), 279-285.

[4] L. Leindler, A new class of numerical sequences and its applications to sine and cosine series, Anal. Math., 28 (2002), 279-286.

[5] L. Leindler, A note on the uniform convergence and boundedness of a new class of sine series, Analysis Math., 31 (2005), 269-275.

[6] L. Leindler, Embedding results regarding strong approximation of sine series, Acta Sci. Math. (Szeged), 71 (1-2) (2005), 91-103.

[7] L. Leindler, A new extension of monotone sequence and its application,, J. Inequal. Pure and Appl. Math., 7(1) (2006), Art. 39, 7 pp.

[8] L. Leindler, Necessary and sufficient conditions for uniform convergence and boundedness of a general class of sine series, Austral. J. Math. Anal. Appl., 4 (1) (2007), Art. 10.

[9] L. Leindler, Integrability conditions pertaining to Orlicz space, J. Inequal. Pure and Appl. Math., 8(2) (2007), Art. 38, 6 pp.

[10] J. R. Nurcombe, On the uniform convergence of sine series with quasimonotone coefficients, J. Math. Anal. Appl., 166 (20) (1992), 577-581.

[11] S. Tikhonov, Trigonometric series with general monotone coefficients, J. Math. Anal. Appl., 326(1) (2007), 721-735.

[12] S. Tikhonov, On uniform convergence of trigonometric series. Mat. Zametki, 81(2) (2007), 304-310, translation in Math. Notes, 81(2) (2007), 268-274.

[13] S. Tikhonov, Best approximation and moduli of smoothness: Computation and equivalence theorems, J. Approx. Theory, 153 (2008), 19-39.

[14] B. Szal, A note on the uniform convergence and boundedness a generalized class of sine series, Commentat. Math., 48 1 (2008), 85-94.

[15] D. S. Yu and S. P. Zhou, A generalization of monotonicity and applications, Acta Math. Hungar., 115 (3) (2007), 247-267.

[16] D. S. Yu, S. P. Zhou and P. Zhou, Ultimate generalization to monotonicity for uniform convergence of trigonometric series, available online at http://arXiv/abs/0704.1865.

[17] D. S. Yu, S. P. Zhou and P. Zhou, On the relations among best approximation and Fourier coefficients, Sci. in China Series A: Math, 51 (2008), 1883-1894.

[18] A. Zygmund, Trigonometric series, Vol. I, University Press (Cambridge, 1959).